

Study of changes of macro- and microelements composition in the cerebrospinal fluid in patients with consequences of craniocerebral trauma

Aliev M.¹, Mamadaliev A.²

Изучение изменений состава макро- и микроэлементов в ликворе у больных с последствиями краниocereбральной травмы

Алиев М. А.¹, Мамадалиев А. М.²

¹Алиев Мансур Абдухоликович / Aliev Mansur Abdukholikovich – старший научный сотрудник - соискатель курса нейрохирургии;

²Мамадалиев Абдурахмон Маматкулович / Mamadaliev Abdurakhmon Matatkulovich – доктор медицинских наук, академик, профессор, зав. курсом нейрохирургии, Академия медико-технических наук РФ,

Самаркандский государственный медицинский институт, г. Самарканд, Узбекистан

Abstract: the aim of this research is to investigate the macro- and microelements composition in the cerebrospinal fluid of patients with different consequences of craniocerebral trauma before and after complex treatment with the use of endolumbal and intracystal introduction of ozone and pyracetam in dynamics. Thus, it may be noted positive changes in the metabolism of macro- and microelements in the cerebrospinal fluid of patients.

Аннотация: изучен состав макро- и микроэлементов в ликворе до и после лечения с использованием эндолумбальной инсуффляции озона и ноотропа и эндокристалльного введения озона при последствиях краниocereбральной травмы. Получены положительные динамические изменения после лечения.

Keywords: macroelements, microelements, cerebrospinal fluid, ozone.

Ключевые слова: макроэлементы, микроэлементы, ликвор, озон.

УДК/UDC-577.17.049:616.8.001

Introduction: Craniocerebral trauma (CCT) for a long time is one of the most actual and complex problem of modern neurosurgery, that many domestic and foreign authors noted in their publications [2, 12, 13, 17, 18]. This is stipulated not only by a high frequency of occurrence, the complexity of the pathogenesis, clinical manifestations and high mortality, but also by an enormous economic damage. One of the most important circumstances stipulating the actuality of the problem is the frequent victims' disability, which arises due to the development of various pathological conditions, persistent symptom complex, united under the name of the consequences of CCT. Just their formation, in the most cases stipulated economic damage caused by cranial injuries [12, 13, 18].

Ozonated saline solution has been successfully used by intravenous introduction in patients with severe CCT in the acute period [1, 4, 5, 11, 16]. One of the first endolumbal introductions of ozone-oxygen mixture has been carried out in 1967, by A. B. Bolgaev in patients with post-traumatic epilepsy. S. D. Madiyarov (1990) has been used once endolumbal introduction of ozone-oxygen mixture as the prevention of cerebral arachnoiditis in patients with severe TBI in the volume of 15 cm³. In 1994, M. K. Agzamov has been carried out scientific researchers on the application of nootropic-ozone mixture in the complex treatment of severe CCT [1, 16]. In 2007, 25 patients with meningitis of different etiologies have been successfully treated by V. M. Belopukhov and his colleagues with the use of endolumbal introduction of ozone-oxygen mixture in the complex treatment [11].

The beginning of the XXI century was marked not only the accumulation of fundamental knowledge in the field of neuroimmunology and neuroelementology, but also the beginning of the application of this knowledge in practice, according to the concept of metabolic protection of the brain [7, 10, 14].

Elaborated in the recent years problem of determining the role and significance of separate chemical elements presenting in tissues and biological fluids of human organism is important both in normal condition and in a variety of diseases, in particular of the nervous system. Variations in the content of macro- and microelements, imbalance of metalloligands' homeostasis caused by both food and ecological factors, as well as somatic diseases which change the status of the nervous system and form an unfavorable background not only for the beginning and development of injures and diseases of the nervous system, but also for the restoration process. The above changes have significantly influence on the understanding of neurorehabilitation measures [7, 8, 9, 10, 19, 20, 21]. The study of domestic and foreign literatures have been showed the absence of researchers devoted to the content of macro- and microelements in the cerebrospinal fluid in the consequences of CCT.

The aim of the study: The aim of this research is to investigate the composition of macro- and microelements in the cerebrospinal fluid of patients with different consequences of CCT before and after complex treatment with the use of endolumbal and intracystal introduction of ozone and pyracetam in dynamics.

Materials and methods: The data of 83 patients with various consequences of TBI in age from 1 year to 60 years old (63 men and 20 women) have been included in the investigation who were hospitalized in neurosurgical clinic of Samarkand Medical Institute from 2009 to 2014. All patients equally with clinical and neurological X-ray investigations (MRI, CT) were carried out laboratory investigations to determine the macro- and microelements (K, Na, Cl, P, Mg, Fe) in the cerebrospinal fluid with the use of «Roche-Hitachi» analyzer before and after 4-6 months treatment. Among the examined patients in 31 patients (37,3 %) it has been diagnosed with post-traumatic cerebral arachnoiditis (PTCA), in 21 patients (25,3 %) it has been diagnosed post-traumatic chronic subdural hematoma (PCSH), in 15 patients (18,1 %) it has been diagnosed post-traumatic epilepsy (PE), in 13 patients (15,7 %) it has been diagnosed post-traumatic arachnoid cyst (PTAC) and in 3 patients (3,6 %) it has been diagnosed chronic vegetative status (CVS) (Figures 1, 2).

For the treatment of patients with consequences of CCT we offered the new methods endocystal introduction of ozone and endolumbal introduction ozone with nootropics (certificates of priority № IAP 2011 0419 and 2011 0148 № IAP).

The method of treatment of arachnoid liquor cysts is that after the imposition of milling holes it has been made dissection and excision of the cyst walls and connection it with subarachnoid and subdural spaces and then vinyl chloride or silicone catheter has been introduced into the cystic cavity and through this catheter with the use of medical syringe it has been injected ozone in the amount of 10-30 cm³ depending on the cyst size, the catheter is left for 3-5 days for the re-introduction of ozone.

The next method is the way endolumbal insufflation of ozone and pyracetam in patients with different consequences of TBI, as mentioned above. These patients under sterile conditions after premedication and local anesthesia by the use of novocain solution 0,5 % – 10,0 ml, a lumbar puncture was performed between the 3rd and 4th lumbar vertebrae, and then it has been evacuated liquor (20-40 ml depending on the liquor pressure) and endolumbal injected first ozone (10-30 cm³), and then pyracetam 2,5–3 % from 200 mg to 1000 mg depending on the age of the patient.

Patients were carried out the following methods of treatment: in patients with PCSH it has been carried out mini-invasive removal of hematomas and endolumbal insufflation of ozone and pyracetam on 7-8 days after surgery; in patients with PTAC it has been carried out mini-invasive cysts emptying and endocystal introduction of ozone in the day of surgery and for 3-4 days after surgery. In PCA, PTE and in patients with CVS after severe CCT it has been conducted endolumbal insufflation ozone and 2-3 % solution of pyracetam (doubly per treatment course).




Figure 1

Forms of consequences of CCT (MRI and CT examples):

- a). Posttraumatic cystic cerebral arachnoidit. CT of patient K. Is determined by a small arachnoid cysts in the left frontal region and cystic-adhesive changes of convexital areas of the brain.*
- b). Posttraumatic epilepsy. MRI of patient R. Determined by the seat of epilepsy - posttraumatic cystic-gliosis in the right frontal region of the brain.*
- c). Posttraumatic arachnoidal cyst. MRI of patient B. Is determined by a arachnoid cysts in the right temporal-basal region of the brain.*
- d). Posttraumatic chronic subdural hematoma. MRI (axial, coronar scans) of patient A. Determined by chronic subdural hematoma in the left hemisphere of the brain.*
- e). Chronic vegetative status. MRI of patient J. Determined by bilateral subdural hydroma fronto-temporal-parietal lobes and post-traumatic atrophic processes of the brain*




Figure 2. Distribution of patients according to the forms of consequences of CCT

Results and discussion: In order to determine the value of pathogenetic changes of macro- and microelements in the brain tissues during the consequences of CCT after using endocystal and endolumbal introduction of ozone and pyracetam, laboratory tests including determination of chemical elements phosphorus (P), calcium (Ca), iron (Fe), magnesium (Mg), sodium (Na), potassium (K) and chlorine (Cl) in the blood serum and in liquor, which play an important role in the activities CNS were performed.

As shown in Table 2 in the liquor tests of these patients it has been observed amounts Ca has been decreased in 89,2 % of patients and amounts of elements such as K and Cl were low in all (100 %) patients. Minimal and maximal amounts of P, Fe and Na in the liquor were normal, but the average amounts of P and Na were high (41 % and 86,7 % accordingly). An increased amount of Mg in the liquor has been noted in 38,6 % (32) of patients in comparison with maximal indexes (Table 2).

Thus, the quantitative changes of all macro- and microelements in the cerebro-spinal fluid in a range of research were noted and once again it has been proved the occurrence of this condition as a result of deep metabolic-biochemical deficiency in the human organism during different consequences of CCT, it gives us the basis to introduce new pathogenetically proved and effective treatment methods in the intermediate and distant periods of traumatic brain disease.

Table 1. Quantitative indexes of macro- and microelements in the liquor in patients with consequences of CCT (before treatment)

Elements	Average indexes (mmol/l)		Minimal indexes (mmol/l)		Maximal indexes (mmol/l)	
	In normal condition	Result	In normal condition	Result	In normal condition	Result
P	0,5	0,83	0,39	0,24	0,68	1,43
Ca	1,4	0,99	1,12	0,69	1,75	1,18
Fe	17	14	5,4	2,1	28,6	25,9
Mg	1,3	1,51	1,23	1,22	1,4	2,06
Na	132,5	140,2	120	121,1	145	151,9
K	3,7	2,3	3,07	2,01	4,35	2,68
Cl	204,5	149,3	197	119,5	212	192

Taking into consideration the above conditions, it has been used the new method of treatment – endolumbal and endocystal insufflation of ozone and pyracetam in the intermediate and distant periods of traumatic disease of the brain and the following results were obtained. On the expiry of 3-6 months after treatment the amount of

macro- and microelements in the cerebrospinal fluid were tested.

Dynamic changes of the structure of macro- and microelements in cerebrospinal fluid after treatment were in the following way: it has been noted the elevated amount of Ca from 0,99 mmol/l before treatment to 1,34 (in 1,4 times) mmol/l; the average index of K has been elevated from 2,3 mmol/l to 3,5 mmol/l (in 1,5 time) and the average index of Cl has been elevated from 149,3 mmol/l to 165,3 mmol/l (in 1,1 time) (Fig. 3).

It has been noted reduction of the average indexes of macro- and microelements which according to the quantitative indexes were up to the normal values: the amount of P was reduced from 0,56 mmol/l to 0,53 mmol/l, the amount of Na was reduced from 140,2 mmol/l to 138,7 mmol/l and the amount of Mg was reduced from 1,51 mmol/l to 1,5 mmol/l (Fig. 4).




Figure 3. Changes in the structure of macro- and microelements in the cerebrospinal fluid in patients with the consequences of the CCT




Figure 4. Changes in the structure of macro- and microelements in the cerebrospinal fluid in patients with the consequences of the CCT

Conclusions:

- Quantitative changes of macro- and microelements in the blood and in cerebrospinal fluid in patients with the consequences of the craniocerebral trauma could cause criterion of the metabolic disorders in the patients' organism and could be the index of the clinical-neurological pathological changes.
- The investigation of the indexes of macro- and microelements in dynamics in patients with the consequences of the craniocerebral trauma could give us the possibility to determine the effectiveness of the pathogenetical treatment of patients.
- Endocystical introduction of ozone and endolumbal insufflation of ozone and introduction of nootropics led to the normalization of macro- and microelements in the blood and in cerebrospinal fluid because of the metabolic improvement in the organism and it could allow us to reach the early restoration of clinical-neurological disorders in patients with the consequences of the craniocerebral trauma.
- Thus, it may be noted positive changes in the metabolism of macro- and microelements in the blood serum and cerebrospinal fluid of patients who were treated according to our suggested methods – endolumbal introductions and nootropic ozone and endocystical introductions of ozone.

References

1. Agzamov M. K. (1994). Comprehensive assessment of the efficacy of nootropic-ozone therapy in severe traumatic brain injury (The dissertation of candidate of medical sciences), Samarkand State Medical Institute, Uzbekistan, in Russian.
2. Akshulakov S. K. (1995). Clinico-epidemiological study of acute traumatic brain injury and its consequences in the Republic of Kazakhstan (The dissertation of doctor of medical sciences), Burdenko Neurosurgery Institute, Russian Federation, in Russian.
3. Aliev M. A., Mamadaliev A. M. (2013). Macronutrient composition of biological media in patients with post-traumatic cerebral arachnoiditis. In Dolgov V. V., Shevchenko O. P. (Eds.), Proceedings of VIII All-Russian Scientific-Practical Conference «Analytical reliability and diagnostic value of laboratory medicine», Journal of Laboratory, № 1/2013 (pp. 23). Moscow, Russian Federation.
4. Aliev M. A., Mamadaliev A.M. (2013). Study of Efficacy of Endocystical Ozonotherapy in the Operative Treatment of Posttraumatic Arachnoidal Cysts. In Hee-Won Jung, Black P. (Eds), Proceedings of XV WFNS Congress, (FA0754)., Seoul, Korea.
5. Bakiev S. S. (2002). The use of ozone in the treatment of patients with fractures of the maxillofacial bones (The dissertation of candidate of medical sciences), Samarkand State Medical Institute, Uzbekistan, in Russian.
6. Bolg'ayev A. B., Madiyarov S. D. (1977). Treatment of chronic arachnoiditis via introduction of ozone in the subarachnoid space. Journal of Neurology and Psychiatry, Moscow, № 2. - 227-231, in Russian.
7. Boldyrev A. A., Eshchenko N. D., Ilyukha V. A., Kaivarainen E. I. (2010). Neurochemistry. Proc. manual for schools. Moscow, in Russian.
8. Bradbury M. B. (1999). Transport of Iron in the Blood-Brain-Cerebrospinal Fluid System. J. Neurochem., Vol. 69. No. 2.
9. Connor J. R., Benkovic S. A. (1992). Iron regulation in the brain: histochemical, biochemical, and molecular considerations. Ann. Neurol. 32 Suppl: S51-61.
10. Druzhinin P. V., Novikov L. F., Lysikov Y. A. (2010). Basics of Nutrition. Clinical management, Moscow, in Russian.
11. Dubrovina Y. A. (2007). Endolumbal introduction of ozone-oxygen mixture in the treatment of traumatic brain injury and its complications: Clinical and neurophysiological evaluation (The dissertation of candidate of medical sciences), Kazan, Russian Federation, in Russian.
12. Konovalov A. N., Likhтерman L. B., Potapov A. A. (1998, 2002). Clinical guidelines for traumatic brain injury. Volume № I and III. Moscow, «Antidor», in Russian.
13. Konovalov A. N., Potapov A. A., Likhтерman L. B., Kornienko V. N., Kravchuk A. D., Okhlopov V. A., Zakharova N. Y., & Yakovlev S. B. (2012). Reconstructive and minimally invasive surgery of outcomes of craniocerebral injury. Moscow, In Russian.
14. Mazurina Y. M. (2005). Violation of calcium metabolism in children during the first three years of life with neurological disorders (The dissertation of candidate of medical sciences), Moscow, Russian Federation, in Russian.
15. Mamadaliev A. M. (1988). Predicting outcomes of traumatic brain injury in the acute period (The dissertation of doctor of medical sciences), Burdenko Neurosurgery Institute, Russian Federation, in Russian.

16. *Mamadaliyev A. M., Agzamov M. K.* (1993). A method for treating severe traumatic brain injury in the acute period (The patent for the invention of the Russian Federation № 5042297/14/023170 on 07/21/93), in Russian.
17. *Mamadaliyev A. M., Aliev M. A.* (2009). The Importance of the Duration Disorders of Consciousness to Prognosis of the Outcome of Cranio-Cerebral Trauma. In Black P. (Eds.), Proceedings of XIV WFNS Congress, Boston, USA.
18. *Mamadaliyev A. M., Shodiev A. S., Aliev M. A., Nabiev A. A.* (2011). By studying the composition of macronutrients are in biological media in post-traumatic cerebral arachnoiditis. In Bersnev V. P. (Eds.), Abstracts of 10th Russian Scientific-Practical Conference of «Polenov reading's», St. Petersburg, Russian Federation, in Russian.
19. *Shakirova E. G.* (2006). Changes of plasma micronutrients in Parkinson's disease and Parkinson's syndrome (The dissertation of candidate of medical sciences), Moscow, Russian Federation, in Russian.
20. *Stepanchenko O. A.* (2005). Peculiarities of stroke when changing glucose in the blood and cerebrospinal fluid (The dissertation of candidate of medical sciences), Moscow, Russian Federation, in Russian.
21. *Ullubiev M. A.* (2008). Changes of plasma micronutrients in ischemic hemispheric stroke (The dissertation of candidate of medical sciences), Moscow, Russian Federation, in Russian.